

- inflammation of liver (hepatitis) or problems with the liver

Very rare side effects include:

- reduced rate of breathing (apnoea)
- reduction in the production of a type of white blood cell making infection more likely (granulocytosis)

If you feel very tired, experience unexpected bruising or bleeding or more infections (e.g. colds and sore throats) than usual please tell your doctor. Your doctor may decide to conduct tests on your blood periodically as a result of these symptoms.

Reporting of side effects

If you get any side effects, talk to your doctor or pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the Yellow card scheme at Website:

www.mhra.gov.uk/yellowcard. By reporting side effects you can help provide more information on the safety of this medicine.

5 How to store Cyclizine Hydrochloride Tablets

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the label and carton. The expiry date refers to the last day of that month. Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

6 Contents of the pack and other information

What Cyclizine Hydrochloride Tablets contain

- The active substance is cyclizine hydrochloride.

Each tablet contains 50mg of the active ingredient, cyclizine hydrochloride,
- The other ingredients are maize starch, lactose, povidone and magnesium stearate.

What Cyclizine Hydrochloride Tablets look like and contents of the pack

Cyclizine Hydrochloride Tablets are circular, white tablets with a breakline on one side and plain on the reverse. These tablets are available in blisters of 1, 10, 28, 30, 40, 50, 84, 100 and 500 tablets. They are also available in a bottle containing 100 tablets.

Not all pack sizes may be marketed.

Marketing Authorisation Holder and Manufacturer

Brown & Burk UK Limited
5, Marryat Close
Hounslow West
Middlesex
TW4 5DQ
UK

This leaflet was last revised in 05/2017.

Package leaflet: Information for the patient
Cyclizine Hydrochloride 50mg Tablets

Cyclizine Hydrochloride

Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor, or pharmacist or nurse.

- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.

- If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. See section 4.

- If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. See section 4.

- If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. See section 4.

- If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. See section 4.

- If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. See section 4.

- If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. See section 4.

- If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. See section 4.

- If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. See section 4.

- If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. See section 4.

- If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. See section 4.

- If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. See section 4.

- If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. See section 4.

- If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. See section 4.

- If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. See section 4.

- If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. See section 4.

- If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. See section 4.

- If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. See section 4.

- If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. See section 4.

- If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. See section 4.

- If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. See section 4.

- If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. See section 4.

- If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. See section 4.

- If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. See section 4.

- If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. See section 4.

- If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. See section 4.

- If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. See section 4.

- If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. See section 4.

- If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. See section 4.

- If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. See section 4.

- If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. See section 4.

- If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. See section 4.

- If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. See section 4.

- If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. See section 4.

- If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. See section 4.

- If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. See section 4.

- If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. See section 4.

- If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. See section 4.

- If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. See section 4.

- If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. See section 4.

- If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. See section 4.

- If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. See section 4.

- If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. See section 4.

- If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. See section 4.

- If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. See section 4.

- If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. See section 4.

- If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. See section 4.

- If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. See section 4.

Cyclizine Hydrochloride Tablets contains the active substance Cyclizine hydrochloride, which belongs to a group of medicines known as antihistamines. Antihistamines help to prevent feelings of sickness (nausea) or being sick (vomiting).

Cyclizine Hydrochloride Tablets may be used by adults and children aged 6 years and over. Cyclizine Hydrochloride Tablets may be used for the following conditions:-

- travel and motion sickness
- post-operative nausea and vomiting, as general anaesthetics can sometimes cause sickness

- nausea associated with cancer treatment (radiotherapy) or other medicines

- treatment of sickness due to inner ear problems, such as Meniere's disease

- treatment of sickness due to inner ear problems, such as Meniere's disease

- treatment of sickness due to inner ear problems, such as Meniere's disease

- treatment of sickness due to inner ear problems, such as Meniere's disease

- treatment of sickness due to inner ear problems, such as Meniere's disease

- treatment of sickness due to inner ear problems, such as Meniere's disease

- treatment of sickness due to inner ear problems, such as Meniere's disease

- treatment of sickness due to inner ear problems, such as Meniere's disease

- treatment of sickness due to inner ear problems, such as Meniere's disease

- treatment of sickness due to inner ear problems, such as Meniere's disease

- treatment of sickness due to inner ear problems, such as Meniere's disease

- treatment of sickness due to inner ear problems, such as Meniere's disease

- treatment of sickness due to inner ear problems, such as Meniere's disease

- treatment of sickness due to inner ear problems, such as Meniere's disease

- treatment of sickness due to inner ear problems, such as Meniere's disease

- treatment of sickness due to inner ear problems, such as Meniere's disease

- treatment of sickness due to inner ear problems, such as Meniere's disease

- treatment of sickness due to inner ear problems, such as Meniere's disease

- treatment of sickness due to inner ear problems, such as Meniere's disease

- treatment of sickness due to inner ear problems, such as Meniere's disease

- treatment of sickness due to inner ear problems, such as Meniere's disease

- treatment of sickness due to inner ear problems, such as Meniere's disease

- treatment of sickness due to inner ear problems, such as Meniere's disease

- treatment of sickness due to inner ear problems, such as Meniere's disease

- treatment of sickness due to inner ear problems, such as Meniere's disease

- treatment of sickness due to inner ear problems, such as Meniere's disease

- treatment of sickness due to inner ear problems, such as Meniere's disease

- treatment of sickness due to inner ear problems, such as Meniere's disease

- treatment of sickness due to inner ear problems, such as Meniere's disease

- treatment of sickness due to inner ear problems, such as Meniere's disease

- treatment of sickness due to inner ear problems, such as Meniere's disease

- treatment of sickness due to inner ear problems, such as Meniere's disease

- treatment of sickness due to inner ear problems, such as Meniere's disease

- treatment of sickness due to inner ear problems, such as Meniere's disease

- treatment of sickness due to inner ear problems, such as Meniere's disease

- treatment of sickness due to inner ear problems, such as Meniere's disease

- treatment of sickness due to inner ear problems, such as Meniere's disease

- treatment of sickness due to inner ear problems, such as Meniere's disease

- treatment of sickness due to inner ear problems, such as Meniere's disease

- treatment of sickness due to inner ear problems, such as Meniere's disease

- treatment of sickness due to inner ear problems, such as Meniere's disease

- treatment of sickness due to inner ear problems, such as Meniere's disease

- treatment of sickness due to inner ear problems, such as Meniere's disease

- treatment of sickness due to inner ear problems, such as Meniere's disease

- treatment of sickness due to inner ear problems, such as Meniere's disease

- treatment of sickness due to inner ear problems, such as Meniere's disease

- treatment of sickness due to inner ear problems, such as Meniere's disease

Warnings and precautions

Talk to your doctor or pharmacist or nurse before taking Cyclizine Hydrochloride Tablets.

Take special care with Cyclizine Hydrochloride Tablets, and tell your doctor or pharmacist if you:

- suffer from an eye disease caused by a rise of pressure within the eye (glaucoma)
- have an obstructive bowel condition

- have any liver problems
- suffer from epilepsy

- are male and have an enlarged prostate gland (difficulty and slowness passing urine)

- have been told your heart is not working properly (heart failure)

- have an inherited disorder which can lead to a sensitivity of sunlight called porphyria

- have been told by your doctor that you have an intolerance to some sugars

- suffer from urinary retention (experience difficulty passing urine)

- suffer from phaeochromocytoma (tumour of the medulla of the adrenal glands)

- suffer from high blood pressure

- suffer from high blood pressure

- suffer from high blood pressure

- suffer from high blood pressure

- suffer from high blood pressure

- suffer from high blood pressure

- suffer from high blood pressure

- suffer from high blood pressure

- suffer from high blood pressure

- suffer from high blood pressure

- suffer from high blood pressure

- suffer from high blood pressure

- suffer from high blood pressure

- suffer from high blood pressure

- suffer from high blood pressure

- suffer from high blood pressure

- suffer from high blood pressure

- suffer from high blood pressure

- suffer from high blood pressure

- suffer from high blood pressure

- suffer from high blood pressure

- suffer from high blood pressure

- suffer from high blood pressure

- suffer from high blood pressure

- suffer from high blood pressure

- suffer from high blood pressure

- suffer from high blood pressure

- suffer from high blood pressure

- suffer from high blood pressure

- suffer from high blood pressure

- suffer from high blood pressure

- suffer from high blood pressure

- suffer from high blood pressure

- suffer from high blood pressure

- suffer from high blood pressure

- suffer from high blood pressure

- suffer from high blood pressure

- suffer from high blood pressure

- suffer from high blood pressure

- suffer from high blood pressure

- suffer from high blood pressure

- suffer from high blood pressure

- any medicine which belong to a group of medicines called anticholinergics

- drugs which can damage the ear (ototoxic) for example aminoglycoside antibacterials.

Cyclizine Hydrochloride Tablets with alcohol

You should avoid taking alcohol whilst you are on Cyclizine Hydrochloride Tablets.

Pregnancy and breast-feeding

If you are pregnant, think you may be pregnant or are planning to have a baby, ask your doctor or pharmacist for advice before taking this medicine.

Do not take these tablets if you are breast-feeding.

Driving and using machines

You can drive while taking Cyclizine Hydrochloride Tablets but do not drive until you know how the tablets affect you.

Cyclizine Hydrochloride Tablets may make you feel dizzy. If they affect you in this way, do not drive or operate any machinery.

Cyclizine Hydrochloride Tablets contain lactose. If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking these tablets.

Cyclizine Hydrochloride Tablets contain lactose. If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking these tablets.

Cyclizine Hydrochloride Tablets contain lactose. If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking these tablets.

Cyclizine Hydrochloride Tablets contain lactose. If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking these tablets.

Cyclizine Hydrochloride Tablets contain lactose. If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking these tablets.

Cyclizine Hydrochloride Tablets contain lactose. If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking these tablets.

Cyclizine Hydrochloride Tablets contain lactose. If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking these tablets.

Cyclizine Hydrochloride Tablets contain lactose. If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking these tablets.

Cyclizine Hydrochloride Tablets contain lactose. If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking these tablets.

Cyclizine Hydrochloride Tablets contain lactose. If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking these tablets.

Cyclizine Hydrochloride Tablets contain lactose. If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking these tablets.

Cyclizine Hydrochloride Tablets contain lactose. If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking these tablets.

Cyclizine Hydrochloride Tablets contain lactose. If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking these tablets.

Cyclizine Hydrochloride Tablets contain lactose. If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking these tablets.

Cyclizine Hydrochloride Tablets contain lactose. If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking these tablets.

Cyclizine Hydrochloride Tablets contain lactose. If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking these tablets.

Cyclizine Hydrochloride Tablets contain lactose. If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking these tablets.

Cyclizine Hydrochloride Tablets contain lactose. If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking these tablets.

Cyclizine Hydrochloride Tablets contain lactose. If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking these tablets.

Cyclizine Hydrochloride Tablets contain lactose. If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking these tablets.

Cyclizine Hydrochloride Tablets contain lactose. If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking these tablets.

Cyclizine Hydrochloride Tablets contain lactose. If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking these tablets.

Cyclizine Hydrochloride Tablets contain lactose. If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking these tablets.

Cyclizine Hydrochloride Tablets contain lactose. If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before